arXiv:1111.3759v1 [math.FA] 16 Nov 2011

Characterizations of operator order for k strictly positive operators [*]

Jian Shi[1] Zongsheng Gao
LMIB & School of Mathematics and Systems Science,
Beihang University, Beijing, 100191, China

Abstract. Let A_i $(i = 1, 2, \cdots, k)$ be bounded linear operators on a Hilbert space. This paper aims to show characterizations of operator order $A_k \geqslant A_{k-1} \geqslant \cdots \geqslant A_2 \geqslant A_1 > 0$ in terms of operator inequalities. Afterwards, an application of the characterizations is given to operator equalities due to Douglas's majorization and factorization theorem.

Keywords and phrases: Positive operator; Further extension of Furuta inequality; Douglas's majorization and factorization theorem.
Mathematics Subject Classification: 47A63.

1 Introduction

A capital letter (such as T) means a bounded linear operator on a Hilbert space $\mathcal{H}$. T is said to be positive (denoted by $T \geqslant 0$) if $(Tx, x) \geqslant 0$ for all $x \in \mathcal{H}$, and T is said to be strictly positive (denoted by $T > 0$) if T is positive and invertible. The usual order $S \geqslant T$ among selfadjoint operators on $\mathcal{H}$ is defined by $(Sx, x) \geqslant (Tx, x)$ for all $x \in \mathcal{H}$. Let I denote the indentity operator.

As an essential and historical extension of the famous Löwner-Heinz inequality: $A \geqslant B \geqslant 0 \Rightarrow A^\alpha \geqslant B^\alpha$ if $\alpha \in [0, 1]$, T. Furuta proved the following operator inequality in 1987.

Theorem 1.1. (Furuta Inequality, [8]) If $A \geqslant B \geqslant 0$, then for each $r \geqslant 0$,

$$(A^{\frac{r}{2}} A^p A^{\frac{r}{2}})^{\frac{1}{q}} \geqslant (A^{\frac{r}{2}} B^p A^{\frac{r}{2}})^{\frac{1}{q}}, \tag{1.1}$$

$$(B^{\frac{r}{2}} A^p B^{\frac{r}{2}})^{\frac{1}{q}} \geqslant (B^{\frac{r}{2}} B^p B^{\frac{r}{2}})^{\frac{1}{q}} \tag{1.2}$$

---

[*] This work is supported by National Natural Science Fund of China (10771011 and 11171013).

[1] Corresponding author.

E-mail addresses: shijian@ss.buaa.edu.cn (J. Shi), zshgao@buaa.edu.cn (Z. Gao).

hold for $p \geqslant 0$ and $q \geqslant 1$ with $(1+r)q \geqslant p+r$.


p
$(1+\boldsymbol{r})\boldsymbol{q} = \boldsymbol{p}+\boldsymbol{r}$
$q=1$
$p=q$
$(1,1)$
$(1,0)$
q
$(0,-r)$

Figure 1: Domain of Furuta inequality

K. Tanahashi showed that the conditions p and q in Figure 1 are best possible for each $r \geqslant 0$. See [16]. It is well known that Furuta inequality has many applications. See [1, 2, 4, 9, 12, 13, 21, 22, 23].

In 1995, T. Furuta showed the following theorem which interpolates Furuta inequality.

Theorem 1.2. (Grand Furuta Inequality, [10]) If $A \geqslant B \geqslant 0$ with $A > 0$, then for each $t \in [0,1]$ and $p \geqslant 1$,

$$A^{1-t+r} \geqslant \{A^{\frac{r}{2}}(A^{-\frac{t}{2}}B^pA^{-\frac{t}{2}})A^{\frac{r}{2}}\}^{\frac{1-t+r}{(p-t)s+r}} \tag{1.3}$$

holds for $s \geqslant 1$ and $r \geqslant t$.

K. Tanahashi proved that the exponent value $\frac{1-t+r}{(p-t)s+r}$ of grand Furuta inequality is the best possible in [17]. Afterwards, the proof was improved by T. Yamazaki and M. Fujii et al., respectively. See [19] and [6].

In 2003, grand Furuta inequality was extended by M. Uchiyama in [18] as follows:

Theorem 1.3. (Extended Grand Furuta Inequality, [18]) If $A \geqslant B \geqslant C \geqslant 0$ with $B > 0$, then for each $t \in [0,1]$ and $p \geqslant 1$,

$$A^{1-t+r} \geqslant A^{\frac{r}{2}}(B^{-\frac{t}{2}}C^pB^{-\frac{t}{2}})^s {A^{\frac{r}{2}}}^{\frac{1-t+r}{(p-t)s+r}} \tag{1.4}$$

holds for $s \geqslant 1$ and $r \geqslant t$.

In 2008, grand Furuta inequality was given another extension in [11] as follows:

Theorem 1.4. (Extension of Furuta Inequality, [11]) If $A \geqslant B \geqslant 0$ with $A > 0$, $t \in [0,1]$

and $p_1, p_2, \ldots, p_{2n} \geqslant 1$ for any natural number n, then the following inequality

$$\begin{aligned} A^{1-t+r} \geqslant \{&A^{\frac{r}{2}}[A^{-\frac{t}{2}} \cdots [A^{-\frac{t}{2}}\{A^{\frac{t}{2}}(A^{-\frac{t}{2}}B^{p_1}A^{-\frac{t}{2}})^{p_2} \\ &A^{\frac{t}{2}}\}^{p_3}A^{-\frac{t}{2}}]^{p_4} \cdots A^{-\frac{t}{2}}]^{p_{2n}}A^{\frac{r}{2}}\}^{\frac{1-t+r}{\phi(2n)-t+r}} \end{aligned} \tag{1.5}$$

holds for $r \geqslant t$, where $\phi(2n) = \{\cdots[\{[(p_1 - t)p_2 + t]p_3 - t\}p_4 + t]p_5 - \cdots - t\}p_{2n} + t$.

In 2010, C. Yang and Y. Wang showed the following theorem which interpolates extended grand Furuta inequality in [20].

Theorem 1.5. (Further Extension of Furuta Inequality, [20]) If $A_{2n+1} \geqslant A_{2n} \geqslant A_{2n-1} \geqslant \cdots \geqslant A_3 \geqslant A_2 \geqslant A_1 \geqslant 0$ with $A_2 > 0$, $t_1, t_2, \ldots, t_{n-1}, t_n \in [0, 1]$ and $p_1, p_2, \ldots, p_{2n-1}, p_{2n} \geqslant 1$ for a natural number n, then the following inequality

$$\begin{aligned} A_{2n+1}^{1-t_n+r} \geqslant \{&A_{2n+1}^{\frac{r}{2}}[A_{2n}^{-\frac{t_n}{2}}\{A_{2n-1}^{\frac{t_{n-1}}{2}} \cdots A_5^{\frac{t_2}{2}}[A_4^{-\frac{t_2}{2}}\{A_3^{\frac{t_1}{2}}(A_2^{-\frac{t_1}{2}}A_1^{p_1}A_2^{-\frac{t_1}{2}})^{p_2} \\ &A_3^{\frac{t_1}{2}}\}^{p_3}A_4^{-\frac{t_2}{2}}]^{p_4}A_5^{\frac{t_2}{2}} \cdots A_{2n-1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}]^{p_{2n}}A_{2n+1}^{\frac{r}{2}}\}^{\frac{1-t_n+r}{\psi[2n]-t_n+r}} \end{aligned} \tag{1.6}$$

holds for $r \geqslant t_n$, where $\psi[2n] = \{\cdots[\{[(p_1 - t_1)p_2 + t_1]p_3 - t_2\}p_4 + t_2]p_5 - \cdots - t_n\}p_{2n} + t_n$.

Recently, some beautiful results on characterizations of operator order have been shown, such as [7], [14] and [15]. C.-S. Lin, by using Furuta inequality, showed the characterizations of operator order for two strictly positive operators in [15]. Afterwards, he and Y. J. Cho, by using extended grand Furuta inequality, showed the characterizations of operator order for three strictly positive operators. The aim of the present paper is to show the characterizations of operator order $A_k \geqslant A_{k-1} \geqslant \cdots \geqslant A_2 \geqslant A_1 > 0$ for any positive integer k in terms of operator inequality via further extension of Furuta inequality. An application of the characterizations is given to operator equalities due to Douglas's majorization and factorization theorem.

2 Main results and proofs

In this section, we show the characterizations of operator order for k strictly positive operators. First, we assume that k is an odd integer ($k = 2n + 1$).

Theorem 2.1. Let $A_1, A_2, A_3, \cdots, A_{2n-1}, A_{2n}, A_{2n+1}$ be strictly positive operators. Then the following two assertions are equivalent.

(I) $A_{2n+1} \geqslant A_{2n} \geqslant A_{2n-1} \geqslant \cdots \geqslant A_3 \geqslant A_2 \geqslant A_1$.

(II) If $t_1, t_2, \cdots, t_n \in [0,1]$, $p_1, p_2, \cdots, p_{2n-1}, p_{2n} \geqslant 1$, $\psi[2n] = \{\cdots[\{[(p_1-t_1)p_2+t_1]p_3-t_2\}p_4+t_2]p_5-\cdots-t_n\}p_{2n}+t_n$, then the following inequalities always hold for $r \geqslant t_n$:

(II.1) $$A_{2n+1}^{r-t_n} \geqslant \Big\{A_{2n+1}^{\frac{r}{2}}\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n-1}^{\frac{t_{n-1}}{2}}\cdots A_5^{\frac{t_2}{2}}\big[A_4^{-\frac{t_2}{2}}\cdot\{A_3^{\frac{t_1}{2}}(A_2^{-\frac{t_1}{2}}A_1^{p_1}A_2^{-\frac{t_1}{2}})^{p_2}A_3^{\frac{t_1}{2}}\}^{p_3}\cdot A_4^{-\frac{t_2}{2}}\big]^{p_4}A_5^{\frac{t_2}{2}}\cdots A_{2n-1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n+1}^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}};$$

(II.2) $$A_{2n+1}^{r-t_n} \geqslant \Big\{A_{2n+1}^{\frac{r}{2}}\Big[A_{2n+1}^{-\frac{t_n}{2}}\{A_{2n}^{\frac{t_{n-1}}{2}}\cdots A_6^{\frac{t_2}{2}}\big[A_5^{-\frac{t_2}{2}}\cdot\{A_4^{\frac{t_1}{2}}(A_3^{-\frac{t_1}{2}}A_2^{p_1}A_3^{-\frac{t_1}{2}})^{p_2}A_4^{\frac{t_1}{2}}\}^{p_3}\cdot A_5^{-\frac{t_2}{2}}\big]^{p_4}A_6^{\frac{t_2}{2}}\cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n+1}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n+1}^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}};$$

(II.3) $$A_{2n+1}^{r-t_n} \geqslant \Big\{A_{2n+1}^{\frac{r}{2}}\Big[A_{2n+1}^{-\frac{t_n}{2}}\{A_{2n+1}^{\frac{t_{n-1}}{2}}\cdots A_7^{\frac{t_2}{2}}\big[A_6^{-\frac{t_2}{2}}\cdot\{A_5^{\frac{t_1}{2}}(A_4^{-\frac{t_1}{2}}A_3^{p_1}A_4^{-\frac{t_1}{2}})^{p_2}A_5^{\frac{t_1}{2}}\}^{p_3}\cdot A_6^{-\frac{t_2}{2}}\big]^{p_4}A_7^{\frac{t_2}{2}}\cdots A_{2n+1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n+1}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n+1}^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}};$$

…………

(II.n) $$A_{2n+1}^{r-t_n} \geqslant \Big\{A_{2n+1}^{\frac{r}{2}}\Big[A_{2n+1}^{-\frac{t_n}{2}}\{A_{2n+1}^{\frac{t_{n-1}}{2}}\cdots A_{n+4}^{\frac{t_2}{2}}\big[A_{n+3}^{-\frac{t_2}{2}}\{A_{n+2}^{\frac{t_1}{2}}(A_{n+1}^{-\frac{t_1}{2}}A_n^{p_1}A_{n+1}^{-\frac{t_1}{2}})^{p_2}A_{n+2}^{\frac{t_1}{2}}\}^{p_3} A_{n+3}^{-\frac{t_2}{2}}\big]^{p_4}A_{n+4}^{\frac{t_2}{2}}\cdots A_{2n+1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n+1}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n+1}^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}};$$

(II.n+1) $$A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_1^{\frac{t_{n-1}}{2}}\cdots A_{n-2}^{\frac{t_2}{2}}\big[A_{n-1}^{-\frac{t_2}{2}}\{A_n^{\frac{t_1}{2}}(A_{n+1}^{-\frac{t_1}{2}}A_{n+2}^{p_1}A_{n+1}^{-\frac{t_1}{2}})^{p_2}A_n^{\frac{t_1}{2}}\}^{p_3} A_{n-1}^{-\frac{t_2}{2}}\big]^{p_4}A_{n-2}^{\frac{t_2}{2}}\cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}};$$

…………

(II.2n-2) $$A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_1^{\frac{t_{n-1}}{2}}\cdots A_{2n-5}^{\frac{t_2}{2}}\big[A_{2n-4}^{-\frac{t_2}{2}}\{A_{2n-3}^{\frac{t_1}{2}}\cdot(A_{2n-2}^{-\frac{t_1}{2}}A_{2n-1}^{p_1}A_{2n-2}^{-\frac{t_1}{2}})^{p_2}\cdot A_{2n-3}^{\frac{t_1}{2}}\}^{p_3}A_{2n-4}^{-\frac{t_2}{2}}\big]^{p_4}A_{2n-5}^{\frac{t_2}{2}}\cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}};$$

(II.2n-1) $$A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_2^{\frac{t_{n-1}}{2}}\cdots A_{2n-4}^{\frac{t_2}{2}}\big[A_{2n-3}^{-\frac{t_2}{2}}\{A_{2n-2}^{\frac{t_1}{2}}\cdot(A_{2n-1}^{-\frac{t_1}{2}}A_{2n}^{p_1}A_{2n-1}^{-\frac{t_1}{2}})^{p_2}\cdot A_{2n-2}^{\frac{t_1}{2}}\}^{p_3}A_{2n-3}^{-\frac{t_2}{2}}\big]^{p_4}A_{2n-4}^{\frac{t_2}{2}}\cdots A_2^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}};$$

(II.2n) $$A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_2^{-\frac{t_n}{2}}\{A_3^{\frac{t_{n-1}}{2}}\cdots A_{2n-3}^{\frac{t_2}{2}}\big[A_{2n-2}^{-\frac{t_2}{2}}\{A_{2n-1}^{\frac{t_1}{2}}\cdot(A_{2n}^{-\frac{t_1}{2}}A_{2n+1}^{p_1}A_{2n}^{-\frac{t_1}{2}})^{p_2}\cdot A_{2n-1}^{\frac{t_1}{2}}\}^{p_3}A_{2n-2}^{-\frac{t_2}{2}}\big]^{p_4}A_{2n-3}^{\frac{t_2}{2}}\cdots A_3^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_2^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}}.$$

Proof. (I)⇒(II) Applying Löwner-Heinz inequality for $\frac{r-t_n}{1-t_n+r}$ to further extension of Furuta inequality, (II.1) is obtained; Replacing $A_1, A_2, A_3, \cdots, A_{2n-1}, A_{2n}$ by $A_2, A_3, A_4, \cdots, A_{2n}, A_{2n+1}$ in (II.1), respectively, (II.2) is obtained; Replacing $A_2, A_3, A_4, \cdots, A_{2n-1}, A_{2n}$ by $A_3, A_4, A_5, \cdots, A_{2n}, A_{2n+1}$ in (II.2), respectively, (II.3) is obtained. Similarly, we can obtain (II.4), (II.5), $\cdots$, (II.n).

If we replace $A_1, A_2, A_3 \cdots, A_{2n-1}, A_{2n}, A_{2n+1}$ by $A_{2n+1}^{-1}, A_{2n}^{-1}, A_{2n-1}^{-1}, \cdots, A_3^{-1}$, A_2^{-1}, A_1^{-1} in (II.1), (II.2), (II.3), $\cdots$, (II.n), respectively, and take reverse, then (II.2n), (II.2n-1), (II.2n-2), $\cdots$, (II.n+1) hold.

(II)$\Rightarrow$(I) Because each A_i is strictly positive and bounded, there exist u_i and v_i such that $+\infty > u_i I \geqslant A_i \geqslant v_i I > 0$ $(i = 1, 2, \cdots, 2n+1)$. If we take $p_1 = p_3 = p_4 = \cdots = p_{2n} = 1$, $t_1 = t_2 = \cdots = t_n = 1$, $r = 2$ in (II.1), then we have

$$\begin{aligned} &A_{2n+1} \\ &\geqslant \{A_{2n+1}A_{2n}^{-\frac{1}{2}}A_{2n-1}^{\frac{1}{2}}\cdots A_5^{\frac{1}{2}}A_4^{-\frac{1}{2}}A_3^{\frac{1}{2}}(A_2^{-\frac{1}{2}}A_1A_2^{-\frac{1}{2}})^{p_2}A_3^{\frac{1}{2}}A_4^{-\frac{1}{2}}A_5^{\frac{1}{2}}\cdots A_{2n-1}^{\frac{1}{2}}A_{2n}^{-\frac{1}{2}}A_{2n+1}\}^{\frac{1}{2}}. \end{aligned} \tag{2.1}$$

According to Theorem 6' in [5]: $X \geqslant Y > 0$ with $sI \geqslant X \geqslant tI > 0 \Rightarrow \frac{(s+t)^2}{4st}X^2 \geqslant Y^2$, we can obtain the following inequality by (2.1) and $u_{2n+1}I \geqslant A_{2n+1} \geqslant v_{2n+1}I > 0$.

$$\begin{aligned} &\frac{(u_{2n+1}+v_{2n+1})^2}{4u_{2n+1}v_{2n+1}}A_{2n+1}^2 \\ &\geqslant A_{2n+1}A_{2n}^{-\frac{1}{2}}A_{2n-1}^{\frac{1}{2}}\cdots A_5^{\frac{1}{2}}A_4^{-\frac{1}{2}}A_3^{\frac{1}{2}}(A_2^{-\frac{1}{2}}A_1A_2^{-\frac{1}{2}})^{p_2}A_3^{\frac{1}{2}}A_4^{-\frac{1}{2}}A_5^{\frac{1}{2}}\cdots A_{2n-1}^{\frac{1}{2}}A_{2n}^{-\frac{1}{2}}A_{2n+1}. \end{aligned} \tag{2.2}$$

Then we have

$$\begin{aligned} &\frac{(u_{2n+1}+v_{2n+1})^2}{4u_{2n+1}v_{2n+1}} \cdot \frac{u_{2n}u_{2n-2}\cdots u_6u_4}{v_{2n-1}v_{2n-3}\cdots v_5v_3}I \\ \geqslant\ &\frac{(u_{2n+1}+v_{2n+1})^2}{4u_{2n+1}v_{2n+1}}A_3^{-\frac{1}{2}}A_4^{\frac{1}{2}}A_5^{-\frac{1}{2}}\cdots A_{2n-1}^{-\frac{1}{2}}A_{2n}A_{2n-1}^{-\frac{1}{2}}\cdots A_5^{-\frac{1}{2}}A_4^{\frac{1}{2}}A_3^{-\frac{1}{2}} \\ \geqslant\ &(A_2^{-\frac{1}{2}}A_1A_2^{-\frac{1}{2}})^{p_2}. \end{aligned} \tag{2.3}$$

Thus,

$$\Big[\frac{(u_{2n+1}+v_{2n+1})^2}{4u_{2n+1}v_{2n+1}} \cdot \frac{u_{2n}u_{2n-2}\cdots u_6u_4}{v_{2n-1}v_{2n-3}\cdots v_5v_3}\Big]^{\frac{1}{p_2}}I \geqslant A_2^{-\frac{1}{2}}A_1A_2^{-\frac{1}{2}} \tag{2.4}$$

holds for any $p_2 \geqslant 1$. $A_2 \geqslant A_1$ is obtained by taking $p_2 \to +\infty$.

Similarly, we can obtain $A_3 \geqslant A_2, A_4 \geqslant A_3, \cdots, A_{n+1} \geqslant A_n$ by (II.2), (II.3), $\cdots$, (II.n), respectively.

By the same setting for (2.2n), the following inequality holds according to Theorem 6 in [5]: $X \geqslant Y > 0$ with $s'I \geqslant Y \geqslant t'I > 0 \Rightarrow \frac{(s'+t')^2}{4s't'}X^2 \geqslant Y^2$.

$$\begin{aligned} &\frac{(u_1+v_1)^2}{4u_1v_1}A_1A_2^{-\frac{1}{2}}A_3^{\frac{1}{2}}\cdots A_{2n-2}^{-\frac{1}{2}}A_{2n-1}^{\frac{1}{2}}(A_{2n}^{-\frac{1}{2}}A_{2n+1}A_{2n}^{-\frac{1}{2}})^{p_2}A_{2n-1}^{\frac{1}{2}}A_{2n-2}^{-\frac{1}{2}}\cdots A_3^{\frac{1}{2}}A_2^{-\frac{1}{2}}A_1 \\ &\geqslant A_1^2. \end{aligned} \tag{2.5}$$

Then we have

$$\begin{aligned}
&A_{2n}^{-\frac{1}{2}}A_{2n+1}A_{2n}^{-\frac{1}{2}}\\
\geqslant{}&\Big[\frac{4u_1v_1}{(u_1+v_1)^2}A_{2n-1}^{-\frac{1}{2}}A_{2n-2}^{\frac{1}{2}}\cdots A_3^{-\frac{1}{2}}A_2A_3^{-\frac{1}{2}}\cdots A_{2n-2}^{\frac{1}{2}}A_{2n-1}^{-\frac{1}{2}}\Big]^{\frac{1}{p_2}}\\
\geqslant{}&\Big[\frac{4u_1v_1}{(u_1+v_1)^2}\cdot\frac{v_2v_4\cdots v_{2n-2}}{u_3u_5\cdots u_{2n-1}}\Big]^{\frac{1}{p_2}}I.
\end{aligned}\tag{2.6}$$

$A_{2n+1}\geqslant A_{2n}$ is obtained by taking $p_2\to+\infty$ in (2.6).

Similarly, we can obtain $A_{2n}\geqslant A_{2n-1}, A_{2n-1}\geqslant A_{2n-2},\cdots,A_{n+2}\geqslant A_{n+1}$ by (II.2n-1), (II.2n-2), $\cdots$, (II.n+1), respectively. □

Remark 2.1. If $n=1$, Theorem 2.1 is the main result of [15].

Next, we assume that k is an even integer ($k=2n$).

Theorem 2.2. *Let $A_1,A_2,A_3,\cdots,A_{2n-1},A_{2n}$ be strictly positive operators. Then the following two assertions are equivalent:*

(I) $A_{2n}\geqslant A_{2n-1}\geqslant\cdots\geqslant A_3\geqslant A_2\geqslant A_1$.

(II) *If $t_1,t_2,\cdots,t_n\in[0,1]$, $p_1,p_2,\cdots,p_{2n-1},p_{2n}\geqslant 1$, $\psi[2n]=\{\cdots[\{[(p_1-t_1)p_2+t_1]p_3-t_2\}p_4+t_2]p_5-\cdots-t_n\}p_{2n}+t_n$, then the following inequalities always hold for $r\geqslant t_n$:*

(II.1) $A_{2n}^{r-t_n}\geqslant\Big\{A_{2n}^{\frac{r}{2}}\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n-1}^{\frac{t_{n-1}}{2}}\cdots A_5^{\frac{t_2}{2}}[A_4^{-\frac{t_2}{2}}\cdot\{A_3^{\frac{t_1}{2}}(A_2^{-\frac{t_1}{2}}A_1^{p_1}A_2^{-\frac{t_1}{2}})^{p_2}A_3^{\frac{t_1}{2}}\}^{p_3}\cdot$
$A_4^{-\frac{t_2}{2}}]^{p_4}A_5^{\frac{t_2}{2}}\cdots A_{2n-1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n}^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}}$;

(II.2) $A_{2n}^{r-t_n}\geqslant\Big\{A_{2n}^{\frac{r}{2}}\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n}^{\frac{t_{n-1}}{2}}\cdots A_6^{\frac{t_2}{2}}[A_5^{-\frac{t_2}{2}}\cdot\{A_4^{\frac{t_1}{2}}(A_3^{-\frac{t_1}{2}}A_2^{p_1}A_3^{-\frac{t_1}{2}})^{p_2}A_4^{\frac{t_1}{2}}\}^{p_3}\cdot$
$A_5^{-\frac{t_2}{2}}]^{p_4}A_6^{\frac{t_2}{2}}\cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n}^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}}$;

(II.3) $A_{2n}^{r-t_n}\geqslant\Big\{A_{2n}^{\frac{r}{2}}\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n}^{\frac{t_{n-1}}{2}}\cdots A_7^{\frac{t_2}{2}}[A_6^{-\frac{t_2}{2}}\cdot\{A_5^{\frac{t_1}{2}}(A_4^{-\frac{t_1}{2}}A_3^{p_1}A_4^{-\frac{t_1}{2}})^{p_2}A_5^{\frac{t_1}{2}}\}^{p_3}\cdot$
$A_6^{-\frac{t_2}{2}}]^{p_4}A_7^{\frac{t_2}{2}}\cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n}^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}}$;

$\cdots\cdots\cdots\cdots$

(II.n) $A_{2n}^{r-t_n}\geqslant\Big\{A_{2n}^{\frac{r}{2}}\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n}^{\frac{t_{n-1}}{2}}\cdots A_{n+4}^{\frac{t_2}{2}}[A_{n+3}^{-\frac{t_2}{2}}\{A_{n+2}^{\frac{t_1}{2}}(A_{n+1}^{-\frac{t_1}{2}}A_n^{p_1}A_{n+1}^{-\frac{t_1}{2}})^{p_2}A_{n+2}^{\frac{t_1}{2}}\}^{p_3}$
$A_{n+3}^{-\frac{t_2}{2}}]^{p_4}A_{n+4}^{\frac{t_2}{2}}\cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n}^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}}$;

(II.n+1) $A_1^{r-t_n}\leqslant\Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_1^{\frac{t_{n-1}}{2}}\cdots A_{n-2}^{\frac{t_2}{2}}[A_{n-1}^{-\frac{t_2}{2}}\{A_n^{\frac{t_1}{2}}(A_{n+1}^{-\frac{t_1}{2}}A_{n+2}^{p_1}A_{n+1}^{-\frac{t_1}{2}})^{p_2}A_n^{\frac{t_1}{2}}\}^{p_3}$
$A_{n-1}^{-\frac{t_2}{2}}]^{p_4}A_{n-2}^{\frac{t_2}{2}}\cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}}$;

. . . .

(II.2n-2) $A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_1^{\frac{t_{n-1}}{2}} \cdots A_{2n-5}^{\frac{t_2}{2}}[A_{2n-4}^{-\frac{t_2}{2}}\{A_{2n-3}^{\frac{t_1}{2}} \cdot (A_{2n-2}^{-\frac{t_1}{2}} A_{2n-1}^{p_1} A_{2n-2}^{-\frac{t_1}{2}})^{p_2} \cdot A_{2n-3}^{\frac{t_1}{2}}\}^{p_3} A_{2n-4}^{-\frac{t_2}{2}}]^{p_4} A_{2n-5}^{\frac{t_2}{2}} \cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}} A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}} A_1^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}}$;

(II.2n-1) $A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_2^{\frac{t_{n-1}}{2}} \cdots A_{2n-4}^{\frac{t_2}{2}}[A_{2n-3}^{-\frac{t_2}{2}}\{A_{2n-2}^{\frac{t_1}{2}} \cdot (A_{2n-1}^{-\frac{t_1}{2}} A_{2n}^{p_1} A_{2n-1}^{-\frac{t_1}{2}})^{p_2} \cdot A_{2n-2}^{\frac{t_1}{2}}\}^{p_3} A_{2n-3}^{-\frac{t_2}{2}}]^{p_4} A_{2n-4}^{\frac{t_2}{2}} \cdots A_2^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}} A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}} A_1^{\frac{r}{2}}\Big\}^{\frac{r-t_n}{\psi[2n]-t_n+r}}$.

Proof. Let $A_{2n+1} = A_{2n}$ in Theorem 2.1. □

Together with Theorem 2.1 and Theorem 2.2, we show the characterizations of operator order $A_k \geqslant A_{k-1} \geqslant \cdots \geqslant A_2 \geqslant A_1 > 0$ for any positive integer k. For example, if $k = 5$, we have the following result:

Proposition 2.1. Let A_1, A_2, A_3, A_4 and A_5 be strictly positive operators. Then $A_5 \geqslant A_4 \geqslant A_3 \geqslant A_2 \geqslant A_1$ if and only if the following four operator inequalities

$$A_5^{r-t_2} \geqslant \Big\{A_5^{\frac{r}{2}}\Big[A_4^{-\frac{t_2}{2}}(A_3^{\frac{t_1}{2}}(A_2^{-\frac{t_1}{2}} A_1^{p_1} A_2^{-\frac{t_1}{2}})^{p_2} A_3^{\frac{t_1}{2}})^{p_3} A_4^{-\frac{t_2}{2}}\Big]^{p_4} A_5^{\frac{r}{2}}\Big\}^{\frac{r-t_2}{\psi[4]-t_2+r}}, \tag{2.7}$$

$$A_5^{r-t_2} \geqslant \Big\{A_5^{\frac{r}{2}}\Big[A_5^{-\frac{t_2}{2}}(A_4^{\frac{t_1}{2}}(A_3^{-\frac{t_1}{2}} A_2^{p_1} A_3^{-\frac{t_1}{2}})^{p_2} A_4^{\frac{t_1}{2}})^{p_3} A_5^{-\frac{t_2}{2}}\Big]^{p_4} A_5^{\frac{r}{2}}\Big\}^{\frac{r-t_2}{\psi[4]-t_2+r}}, \tag{2.8}$$

$$A_1^{r-t_2} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_2}{2}}(A_2^{\frac{t_1}{2}}(A_3^{-\frac{t_1}{2}} A_4^{p_1} A_3^{-\frac{t_1}{2}})^{p_2} A_2^{\frac{t_1}{2}})^{p_3} A_1^{-\frac{t_2}{2}}\Big]^{p_4} A_1^{\frac{r}{2}}\Big\}^{\frac{r-t_2}{\psi[4]-t_2+r}}, \tag{2.9}$$

$$A_1^{r-t_2} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_2^{-\frac{t_2}{2}}(A_3^{\frac{t_1}{2}}(A_4^{-\frac{t_1}{2}} A_5^{p_1} A_4^{-\frac{t_1}{2}})^{p_2} A_3^{\frac{t_1}{2}})^{p_3} A_2^{-\frac{t_2}{2}}\Big]^{p_4} A_1^{\frac{r}{2}}\Big\}^{\frac{r-t_2}{\psi[4]-t_2+r}} \tag{2.10}$$

always hold for $p_1, p_2, p_3, p_4 \geqslant 1$, $t_1, t_2 \in [0,1]$ and $r \geqslant t_2$, where $\psi[4] = \{[(p_1 - t_1)p_2 + t_1]p_3 - t_2\}p_4 + t_2$.

Remark 2.2. It should be mentioned that we can not obtain $A_5 \geqslant A_4 \geqslant A_3 \geqslant A_2 \geqslant A_1$ only by (2.7) and (2.10). If $A_1 = \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{u} \end{pmatrix}$, $A_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, $A_3 = \begin{pmatrix} 1 & 0 \\ 0 & u \end{pmatrix}$, $A_4 = \begin{pmatrix} u & 0 \\ 0 & 1 \end{pmatrix}$, $A_5 = \begin{pmatrix} u+\varepsilon & 0 \\ 0 & 1 \end{pmatrix}$, where $u > 1$ and $\varepsilon > 0$, then the five strictly positive operators satisfy (2.7) and (2.10) without satisfying $A_4 \geqslant A_3$.

3 An application

In what follows we give an application of the characterizations in Theorem 2.1 and Theorem 2.2 to operator equalities.

Theorem 3.1. If $A_1, A_2, A_3, \cdots, A_{2n-1}, A_{2n}, A_{2n+1}$ are strictly positive operators, $t_1, t_2, \cdots, t_n \in [0,1]$, $p_1, p_2, \cdots, p_{2n} \geqslant 1$, $\psi[2n] = \{\cdots[\{[(p_1 - t_1)p_2 + t_1]p_3 - t_2\}p_4 + t_2]p_5 - \cdots - t_n\}p_{2n} + t_n$, $r \geqslant t_n$, m is a positive integer such that $(r - t_n)m = \psi[2n] - t_n + r$ with $m \geqslant 2$, then the following assertions are mutually equivalent:

(I) $A_{2n+1} \geqslant A_{2n} \geqslant A_{2n-1} \geqslant \cdots \geqslant A_3 \geqslant A_2 \geqslant A_1$.

(II) The following operator inequalities hold:

(II.1) $A_{2n+1}^{r-t_n} \geqslant \Big\{A_{2n+1}^{\frac{r}{2}}\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n-1}^{\frac{t_{n-1}}{2}} \cdots A_5^{\frac{t_2}{2}}\,[A_4^{-\frac{t_2}{2}} \cdot \{A_3^{\frac{t_1}{2}}(A_2^{-\frac{t_1}{2}}A_1^{p_1}A_2^{-\frac{t_1}{2}})^{p_2}A_3^{\frac{t_1}{2}}\}^{p_3} \cdot$
$A_4^{-\frac{t_2}{2}}]^{p_4}A_5^{\frac{t_2}{2}} \cdots A_{2n-1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n+1}^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

(II.2) $A_{2n+1}^{r-t_n} \geqslant \Big\{A_{2n+1}^{\frac{r}{2}}\Big[A_{2n+1}^{-\frac{t_n}{2}}\{A_{2n}^{\frac{t_{n-1}}{2}} \cdots A_6^{\frac{t_2}{2}}\,[A_5^{-\frac{t_2}{2}} \cdot \{A_4^{\frac{t_1}{2}}(A_3^{-\frac{t_1}{2}}A_2^{p_1}A_3^{-\frac{t_1}{2}})^{p_2}A_4^{\frac{t_1}{2}}\}^{p_3} \cdot$
$A_5^{-\frac{t_2}{2}}]^{p_4}A_6^{\frac{t_2}{2}} \cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n+1}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n+1}^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

(II.3) $A_{2n+1}^{r-t_n} \geqslant \Big\{A_{2n+1}^{\frac{r}{2}}\Big[A_{2n+1}^{-\frac{t_n}{2}}\{A_{2n+1}^{\frac{t_{n-1}}{2}} \cdots A_7^{\frac{t_2}{2}}\,[A_6^{-\frac{t_2}{2}} \cdot \{A_5^{\frac{t_1}{2}}(A_4^{-\frac{t_1}{2}}A_3^{p_1}A_4^{-\frac{t_1}{2}})^{p_2}A_5^{\frac{t_1}{2}}\}^{p_3} \cdot$
$A_6^{-\frac{t_2}{2}}]^{p_4}A_7^{\frac{t_2}{2}} \cdots A_{2n+1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n+1}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n+1}^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

$\cdots\cdots\cdots\cdots$

(II.n) $A_{2n+1}^{r-t_n} \geqslant \Big\{A_{2n+1}^{\frac{r}{2}}\Big[A_{2n+1}^{-\frac{t_n}{2}}\{A_{2n+1}^{\frac{t_{n-1}}{2}} \cdots A_{n+4}^{\frac{t_2}{2}}[A_{n+3}^{-\frac{t_2}{2}}\{A_{n+2}^{\frac{t_1}{2}}(A_{n+1}^{-\frac{t_1}{2}}A_n^{p_1}A_{n+1}^{-\frac{t_1}{2}})^{p_2}A_{n+2}^{\frac{t_1}{2}}\}^{p_3}$
$A_{n+3}^{-\frac{t_2}{2}}]^{p_4}A_{n+4}^{\frac{t_2}{2}} \cdots A_{2n+1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n+1}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n+1}^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

(II.n+1) $A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_1^{\frac{t_{n-1}}{2}} \cdots A_{n-2}^{\frac{t_2}{2}}[A_{n-1}^{-\frac{t_2}{2}}\{A_n^{\frac{t_1}{2}}(A_{n+1}^{-\frac{t_1}{2}}A_{n+2}^{p_1}A_{n+1}^{-\frac{t_1}{2}})^{p_2}A_n^{\frac{t_1}{2}}\}^{p_3}$
$A_{n-1}^{-\frac{t_2}{2}}]^{p_4}A_{n-2}^{\frac{t_2}{2}} \cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

$\cdots\cdots\cdots\cdots$

(II.2n-2) $A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_1^{\frac{t_{n-1}}{2}} \cdots A_{2n-5}^{\frac{t_2}{2}}[A_{2n-4}^{-\frac{t_2}{2}}\{A_{2n-3}^{\frac{t_1}{2}} \cdot (A_{2n-2}^{-\frac{t_1}{2}}A_{2n-1}^{p_1}A_{2n-2}^{-\frac{t_1}{2}})^{p_2} \cdot$
$A_{2n-3}^{\frac{t_1}{2}}\}^{p_3}A_{2n-4}^{-\frac{t_2}{2}}]^{p_4}A_{2n-5}^{\frac{t_2}{2}} \cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

(II.2n-1) $A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_2^{\frac{t_{n-1}}{2}} \cdots A_{2n-4}^{\frac{t_2}{2}}[A_{2n-3}^{-\frac{t_2}{2}}\{A_{2n-2}^{\frac{t_1}{2}} \cdot (A_{2n-1}^{-\frac{t_1}{2}}A_{2n}^{p_1}A_{2n-1}^{-\frac{t_1}{2}})^{p_2} \cdot$
$A_{2n-2}^{\frac{t_1}{2}}\}^{p_3}A_{2n-3}^{-\frac{t_2}{2}}]^{p_4}A_{2n-4}^{\frac{t_2}{2}} \cdots A_2^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

(II.2n) $A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_2^{-\frac{t_n}{2}}\{A_3^{\frac{t_{n-1}}{2}} \cdots A_{2n-3}^{\frac{t_2}{2}}[A_{2n-2}^{-\frac{t_2}{2}}\{A_{2n-1}^{\frac{t_1}{2}} \cdot (A_{2n}^{-\frac{t_1}{2}}A_{2n+1}^{p_1}A_{2n}^{-\frac{t_1}{2}})^{p_2} \cdot$
$A_{2n-1}^{\frac{t_1}{2}}\}^{p_3}A_{2n-2}^{-\frac{t_2}{2}}]^{p_4}A_{2n-3}^{\frac{t_2}{2}} \cdots A_3^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_2^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$.

(III) There exists strictly positive operators $S_1, S_2, S_3, \cdots, S_{2n-2}, S_{2n-1}, S_{2n}$ satisfying the following operator equalities, respectively, where each S_i $(i = 1, 2, \cdots, 2n)$ is unique with $\|S_i\| \leqslant 1$.

(III.1) $A_{2n+1}^{-\frac{t_n}{2}}S_1(A_{2n+1}^{r-t_n}S_1)^{m-1}A_{2n+1}^{-\frac{t_n}{2}} = A_{2n+1}^{-\frac{t_n}{2}}(S_1A_{2n+1}^{r-t_n})^{m-1}S_1A_{2n+1}^{-\frac{t_n}{2}} =$

$$\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n-1}^{\frac{t_{n-1}}{2}}\cdots A_5^{\frac{t_2}{2}}\big[A_4^{-\frac{t_2}{2}}\{A_3^{\frac{t_1}{2}}(A_2^{-\frac{t_1}{2}}A_1^{p_1}A_2^{-\frac{t_1}{2}})^{p_2}A_3^{\frac{t_1}{2}}\}^{p_3}A_4^{-\frac{t_2}{2}}\big]^{p_4}A_5^{\frac{t_2}{2}}\cdots A_{2n-1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}};$$

(III.2) $A_{2n+1}^{-\frac{t_n}{2}}S_2(A_{2n+1}^{r-t_n}S_2)^{m-1}A_{2n+1}^{-\frac{t_n}{2}} = A_{2n+1}^{-\frac{t_n}{2}}(S_2A_{2n+1}^{r-t_n})^{m-1}S_2A_{2n+1}^{-\frac{t_n}{2}} =$

$$\Big[A_{2n+1}^{-\frac{t_n}{2}}\{A_{2n}^{\frac{t_{n-1}}{2}}\cdots A_6^{\frac{t_2}{2}}\big[A_5^{-\frac{t_2}{2}}\{A_4^{\frac{t_1}{2}}(A_3^{-\frac{t_1}{2}}A_2^{p_1}A_3^{-\frac{t_1}{2}})^{p_2}A_4^{\frac{t_1}{2}}\}^{p_3}A_5^{-\frac{t_2}{2}}\big]^{p_4}A_6^{\frac{t_2}{2}}\cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n+1}^{-\frac{t_n}{2}}\Big]^{p_{2n}};$$

(III.3) $A_{2n+1}^{-\frac{t_n}{2}}S_3(A_{2n+1}^{r-t_n}S_3)^{m-1}A_{2n+1}^{-\frac{t_n}{2}} = A_{2n+1}^{-\frac{t_n}{2}}(S_3A_{2n+1}^{r-t_n})^{m-1}S_3A_{2n+1}^{-\frac{t_n}{2}} =$

$$\Big[A_{2n+1}^{-\frac{t_n}{2}}\{A_{2n+1}^{\frac{t_{n-1}}{2}}\cdots A_7^{\frac{t_2}{2}}\big[A_6^{-\frac{t_2}{2}}\{A_5^{\frac{t_1}{2}}(A_4^{-\frac{t_1}{2}}A_3^{p_1}A_4^{-\frac{t_1}{2}})^{p_2}A_5^{\frac{t_1}{2}}\}^{p_3}A_6^{-\frac{t_2}{2}}\big]^{p_4}A_7^{\frac{t_2}{2}}\cdots A_{2n+1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n+1}^{-\frac{t_n}{2}}\Big]^{p_{2n}};$$

$\cdots\cdots\cdots\cdots$

(III.n) $A_{2n+1}^{-\frac{t_n}{2}}S_n(A_{2n+1}^{r-t_n}S_n)^{m-1}A_{2n+1}^{-\frac{t_n}{2}} = A_{2n+1}^{-\frac{t_n}{2}}(S_nA_{2n+1}^{r-t_n})^{m-1}S_nA_{2n+1}^{-\frac{t_n}{2}} =$

$$\Big[A_{2n+1}^{-\frac{t_n}{2}}\{A_{2n+1}^{\frac{t_{n-1}}{2}}\cdots A_{n+4}^{\frac{t_2}{2}}\big[A_{n+3}^{-\frac{t_2}{2}}\{A_{n+2}^{\frac{t_1}{2}}(A_{n+1}^{-\frac{t_1}{2}}A_n^{p_1}A_{n+1}^{-\frac{t_1}{2}})^{p_2}A_{n+2}^{\frac{t_1}{2}}\}^{p_3}A_{n+3}^{-\frac{t_2}{2}}\big]^{p_4}A_{n+4}^{\frac{t_2}{2}}\cdots A_{2n+1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n+1}^{-\frac{t_n}{2}}\Big]^{p_{2n}};$$

(III.n+1) $A_1^{-\frac{t_n}{2}}S_{n+1}^{-1}(A_1^{r-t_n}S_{n+1}^{-1})^{m-1}A_1^{-\frac{t_n}{2}} = A_1^{-\frac{t_n}{2}}(S_{n+1}^{-1}A_1^{r-t_n})^{m-1}S_{n+1}^{-1}A_1^{-\frac{t_n}{2}} =$

$$\Big[A_1^{-\frac{t_n}{2}}\{A_1^{\frac{t_{n-1}}{2}}\cdots A_{n-2}^{\frac{t_2}{2}}\big[A_{n-1}^{-\frac{t_2}{2}}\{A_n^{\frac{t_1}{2}}(A_{n+1}^{-\frac{t_1}{2}}A_{n+2}^{p_1}A_{n+1}^{-\frac{t_1}{2}})^{p_2}A_n^{\frac{t_1}{2}}\}^{p_3}A_{n-1}^{-\frac{t_2}{2}}\big]^{p_4}A_{n-2}^{\frac{t_2}{2}}\cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}};$$

$\cdots\cdots\cdots\cdots$

(III.2n-2) $A_1^{-\frac{t_n}{2}}S_{2n-2}^{-1}(A_1^{r-t_n}S_{2n-2}^{-1})^{m-1}A_1^{-\frac{t_n}{2}} = A_1^{-\frac{t_n}{2}}(S_{2n-2}^{-1}A_1^{r-t_n})^{m-1}S_{2n-2}^{-1}A_1^{-\frac{t_n}{2}} =$

$$\Big[A_1^{-\frac{t_n}{2}}\{A_1^{\frac{t_{n-1}}{2}}\cdots\big[A_{2n-4}^{-\frac{t_2}{2}}\{A_{2n-3}^{\frac{t_1}{2}}(A_{2n-2}^{-\frac{t_1}{2}}A_{2n-1}^{p_1}A_{2n-2}^{-\frac{t_1}{2}})^{p_2}A_{2n-3}^{\frac{t_1}{2}}\}^{p_3}A_{2n-4}^{-\frac{t_2}{2}}\big]^{p_4}\cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}};$$

(III.2n-1) $A_1^{-\frac{t_n}{2}}S_{2n-1}^{-1}(A_1^{r-t_n}S_{2n-1}^{-1})^{m-1}A_1^{-\frac{t_n}{2}} = A_1^{-\frac{t_n}{2}}(S_{2n-1}^{-1}A_1^{r-t_n})^{m-1}S_{2n-1}^{-1}A_1^{-\frac{t_n}{2}} =$

$$\Big[A_1^{-\frac{t_n}{2}}\{A_2^{\frac{t_{n-1}}{2}}\cdots\big[A_{2n-3}^{-\frac{t_2}{2}}\{A_{2n-2}^{\frac{t_1}{2}}(A_{2n-1}^{-\frac{t_1}{2}}A_{2n}^{p_1}A_{2n-1}^{-\frac{t_1}{2}})^{p_2}A_{2n-2}^{\frac{t_1}{2}}\}^{p_3}A_{2n-3}^{-\frac{t_2}{2}}\big]^{p_4}\cdots A_2^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}};$$

(III.2n) $A_1^{-\frac{t_n}{2}}S_{2n}^{-1}(A_1^{r-t_n}S_{2n}^{-1})^{m-1}A_1^{-\frac{t_n}{2}} = A_1^{-\frac{t_n}{2}}(S_{2n}^{-1}A_1^{r-t_n})^{m-1}S_{2n}^{-1}A_1^{-\frac{t_n}{2}} =$

$$\Big[A_2^{-\frac{t_n}{2}}\{A_3^{\frac{t_{n-1}}{2}}\cdots\big[A_{2n-2}^{-\frac{t_2}{2}}\{A_{2n-1}^{\frac{t_1}{2}}(A_{2n}^{-\frac{t_1}{2}}A_{2n+1}^{p_1}A_{2n}^{-\frac{t_1}{2}})^{p_2}A_{2n-1}^{\frac{t_1}{2}}\}^{p_3}A_{2n-2}^{-\frac{t_2}{2}}\big]^{p_4}\cdots A_3^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_2^{-\frac{t_n}{2}}\Big]^{p_{2n}}.$$

Proof. Because (II)⇔(III) holds obviously by Theorem 2.1, we only need to prove that (II)⇔(III).

Firstly, let us prove that (II.1)⇒(III.1). We recall Douglas's majorization and factorization theorem in [3]: $SS^* \leqslant \lambda^2 TT^* \Leftrightarrow$ there exists an operator Q s.t. $TQ = S$, where $\|Q\|^2 = inf\{\mu : SS^* \leqslant \mu TT^*\}$.

By (II.1), there exists an operator E_1 with $\|E_1\| \leqslant 1$ such that

$$\begin{aligned} & A_{2n+1}^{\frac{r-t_n}{2}} E_1 = E_1^* A_{2n+1}^{\frac{r-t_n}{2}} \\ = & \Big\{ A_{2n+1}^{\frac{r}{2}} \Big[A_{2n}^{-\frac{t_n}{2}} \{ A_{2n-1}^{\frac{t_{n-1}}{2}} \cdots \{ A_3^{\frac{t_1}{2}} (A_2^{-\frac{t_1}{2}} A_1^{p_1} A_2^{-\frac{t_1}{2}})^{p_2} A_3^{\frac{t_1}{2}} \}^{p_3} \cdots A_{2n-1}^{\frac{t_{n-1}}{2}} \}^{p_{2n-1}} A_{2n}^{-\frac{t_n}{2}} \Big]^{p_{2n}} A_{2n+1}^{\frac{r}{2}} \Big\}^{\frac{1}{2m}}. \end{aligned} \tag{3.1}$$

Taking $S_1 = E_1 E_1^*$, we have

$$\begin{aligned} & A_{2n+1}^{\frac{r-t_n}{2}} S_1 A_{2n+1}^{\frac{r-t_n}{2}} \\ = & \Big\{ A_{2n+1}^{\frac{r}{2}} \Big[A_{2n}^{-\frac{t_n}{2}} \{ A_{2n-1}^{\frac{t_{n-1}}{2}} \cdots \{ A_3^{\frac{t_1}{2}} (A_2^{-\frac{t_1}{2}} A_1^{p_1} A_2^{-\frac{t_1}{2}})^{p_2} A_3^{\frac{t_1}{2}} \}^{p_3} \cdots A_{2n-1}^{\frac{t_{n-1}}{2}} \}^{p_{2n-1}} A_{2n}^{-\frac{t_n}{2}} \Big]^{p_{2n}} A_{2n+1}^{\frac{r}{2}} \Big\}^{\frac{1}{m}}. \end{aligned} \tag{3.2}$$

According to (3.2) and $S_1 = E_1 E_1^*$, S_1 is unique and strictly positive with $\|S_1\| \leqslant 1$. (3.2) also implies that

$$\begin{aligned} & (A_{2n+1}^{\frac{r-t_n}{2}} S_1 A_{2n+1}^{\frac{r-t_n}{2}})^m = A_{2n+1}^{\frac{r-t_n}{2}} S_1 (A_{2n+1}^{r-t_n} S_1)^{m-1} A_{2n+1}^{\frac{r-t_n}{2}} = A_{2n+1}^{\frac{r-t_n}{2}} (S_1 A_{2n+1}^{r-t_n})^{m-1} S_1 A_{2n+1}^{\frac{r-t_n}{2}} \\ = & A_{2n+1}^{\frac{r}{2}} \Big[A_{2n}^{-\frac{t_n}{2}} \{ A_{2n-1}^{\frac{t_{n-1}}{2}} \cdots \{ A_3^{\frac{t_1}{2}} (A_2^{-\frac{t_1}{2}} A_1^{p_1} A_2^{-\frac{t_1}{2}})^{p_2} A_3^{\frac{t_1}{2}} \}^{p_3} \cdots A_{2n-1}^{\frac{t_{n-1}}{2}} \}^{p_{2n-1}} A_{2n}^{-\frac{t_n}{2}} \Big]^{p_{2n}} A_{2n+1}^{\frac{r}{2}}. \end{aligned} \tag{3.3}$$

Then (III.1) holds by (3.3).

Secondly, we prove that (III.1)⇒ (II.1). By (III.1),

$$\begin{aligned} & \Big\{ A_{2n+1}^{\frac{r}{2}} \Big[A_{2n}^{-\frac{t_n}{2}} \{ A_{2n-1}^{\frac{t_{n-1}}{2}} \cdots \{ A_3^{\frac{t_1}{2}} (A_2^{-\frac{t_1}{2}} A_1^{p_1} A_2^{-\frac{t_1}{2}})^{p_2} A_3^{\frac{t_1}{2}} \}^{p_3} \cdots A_{2n-1}^{\frac{t_{n-1}}{2}} \}^{p_{2n-1}} A_{2n}^{-\frac{t_n}{2}} \Big]^{p_{2n}} A_{2n+1}^{\frac{r}{2}} \Big\}^{\frac{1}{m}} \\ = & \{ A_{2n+1}^{\frac{r-t_n}{2}} S_1 (A_{2n+1}^{r-t_n} S_1)^{m-1} A_{2n+1}^{\frac{r-t_n}{2}} \}^{\frac{1}{m}} \\ = & \{ (A_{2n+1}^{\frac{r-t_n}{2}} S_1 A_{2n+1}^{\frac{r-t_n}{2}}) \cdot (A_{2n+1}^{\frac{r-t_n}{2}} S_1 A_{2n+1}^{\frac{r-t_n}{2}}) \cdots (A_{2n+1}^{\frac{r-t_n}{2}} S_1 A_{2n+1}^{\frac{r-t_n}{2}}) \}^{\frac{1}{m}} \\ = & A_{2n+1}^{\frac{r-t_n}{2}} S_1 A_{2n+1}^{\frac{r-t_n}{2}} \\ \leqslant & A_{2n+1}^{r-t_n}. \end{aligned}$$

The inequality follows form the fact that $S_1 \leqslant \|S_1\| I \leqslant I$, and then (II.1) holds.

By using the same method above, we can prove that (II.2)⇔ (III.2), (II.3)⇔ (III.3), $\cdots$, (II.n)⇔ (III.n), respectively.

Next, we show that (II.2n)⇔(III.2n). Notice that for two strictly positive operators S and T, $S \geqslant T$ if and only if $T^{-1} \geqslant S^{-1}$. Then (II.2n) is equivalent to

$$\begin{aligned} A_1^{-(r-t_n)} \geqslant \Big\{ & A_1^{-\frac{r}{2}} \Big[A_2^{-\frac{t_n}{2}} \{ A_3^{\frac{t_{n-1}}{2}} \cdots A_{2n-3}^{\frac{t_2}{2}} \big[A_{2n-2}^{-\frac{t_2}{2}} \{ A_{2n-1}^{\frac{t_1}{2}} (A_{2n}^{-\frac{t_1}{2}} A_{2n+1}^{p_1} A_{2n}^{-\frac{t_1}{2}})^{p_2} \\ & A_{2n-1}^{\frac{t_1}{2}} \}^{p_3} A_{2n-2}^{-\frac{t_2}{2}} \big]^{p_4} A_{2n-3}^{\frac{t_2}{2}} \cdots A_3^{\frac{t_{n-1}}{2}} \}^{p_{2n-1}} A_2^{-\frac{t_n}{2}} \Big]^{-p_{2n}} A_1^{-\frac{r}{2}} \Big\}^{\frac{1}{m}}. \end{aligned} \tag{3.4}$$

The proof that (3.4)⇔(III.2n) is similar to the proof of that (II.1)⇔(III.1), so we omit it here.

Repeat the method above, we can prove that (II.2n-1)⇔(III.2n-1), (II.2n-2)⇔(III.2n-2), $\cdots$, (II.n+1)⇔(III.n+1). □

Theorem 3.2. If $A_1, A_2, A_3, \cdots, A_{2n-1}, A_{2n}$ are strictly positive operators, $t_1, t_2, \cdots, t_n \in [0,1]$, $p_1, p_2, \cdots, p_{2n} \geqslant 1$, $\psi[2n] = \{\cdots[\{[(p_1-t_1)p_2+t_1]p_3-t_2\}p_4+t_2]p_5-\cdots-t_n\}p_{2n}+t_n$, $r \geqslant t_n$, m is a positive integer such that $(r-t_n)m = \psi[2n]-t_n+r$ with $m \geqslant 2$, then the following assertions are mutually equivalent:

(I) $A_{2n} \geqslant A_{2n-1} \geqslant \cdots \geqslant A_3 \geqslant A_2 \geqslant A_1$.

(II) The following operator inequalities hold:

(II.1) $A_{2n}^{r-t_n} \geqslant \Big\{A_{2n}^{\frac{r}{2}}\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n-1}^{\frac{t_{n-1}}{2}}\cdots A_5^{\frac{t_2}{2}}[A_4^{-\frac{t_2}{2}}\cdot\{A_3^{\frac{t_1}{2}}(A_2^{-\frac{t_1}{2}}A_1^{p_1}A_2^{-\frac{t_1}{2}})^{p_2}A_3^{\frac{t_1}{2}}\}^{p_3}\cdot A_4^{-\frac{t_2}{2}}]^{p_4}A_5^{\frac{t_2}{2}}\cdots A_{2n-1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n}^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

(II.2) $A_{2n}^{r-t_n} \geqslant \Big\{A_{2n}^{\frac{r}{2}}\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n}^{\frac{t_{n-1}}{2}}\cdots A_6^{\frac{t_2}{2}}[A_5^{-\frac{t_2}{2}}\cdot\{A_4^{\frac{t_1}{2}}(A_3^{-\frac{t_1}{2}}A_2^{p_1}A_3^{-\frac{t_1}{2}})^{p_2}A_4^{\frac{t_1}{2}}\}^{p_3}\cdot A_5^{-\frac{t_2}{2}}]^{p_4}A_6^{\frac{t_2}{2}}\cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n}^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

(II.3) $A_{2n}^{r-t_n} \geqslant \Big\{A_{2n}^{\frac{r}{2}}\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n}^{\frac{t_{n-1}}{2}}\cdots A_7^{\frac{t_2}{2}}[A_6^{-\frac{t_2}{2}}\cdot\{A_5^{\frac{t_1}{2}}(A_4^{-\frac{t_1}{2}}A_3^{p_1}A_4^{-\frac{t_1}{2}})^{p_2}A_5^{\frac{t_1}{2}}\}^{p_3}\cdot A_6^{-\frac{t_2}{2}}]^{p_4}A_7^{\frac{t_2}{2}}\cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n}^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

$\cdots\cdots\cdots\cdots$

(II.n) $A_{2n}^{r-t_n} \geqslant \Big\{A_{2n}^{\frac{r}{2}}\Big[A_{2n}^{-\frac{t_n}{2}}\{A_{2n}^{\frac{t_{n-1}}{2}}\cdots A_{n+4}^{\frac{t_2}{2}}[A_{n+3}^{-\frac{t_2}{2}}\{A_{n+2}^{\frac{t_1}{2}}(A_{n+1}^{-\frac{t_1}{2}}A_n^{p_1}A_{n+1}^{-\frac{t_1}{2}})^{p_2}A_{n+2}^{\frac{t_1}{2}}\}^{p_3} A_{n+3}^{-\frac{t_2}{2}}]^{p_4}A_{n+4}^{\frac{t_2}{2}}\cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_{2n}^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

(II.n+1) $A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_1^{\frac{t_{n-1}}{2}}\cdots A_{n-2}^{\frac{t_2}{2}}[A_{n-1}^{-\frac{t_2}{2}}\{A_n^{\frac{t_1}{2}}(A_{n+1}^{-\frac{t_1}{2}}A_{n+2}^{p_1}A_{n+1}^{-\frac{t_1}{2}})^{p_2}A_n^{\frac{t_1}{2}}\}^{p_3} A_{n-1}^{-\frac{t_2}{2}}]^{p_4}A_{n-2}^{\frac{t_2}{2}}\cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

$\cdots\cdots\cdots\cdots$

(II.2n-2) $A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_1^{\frac{t_{n-1}}{2}}\cdots A_{2n-5}^{\frac{t_2}{2}}[A_{2n-4}^{-\frac{t_2}{2}}\{A_{2n-3}^{\frac{t_1}{2}}\cdot(A_{2n-2}^{-\frac{t_1}{2}}A_{2n-1}^{p_1}A_{2n-2}^{-\frac{t_1}{2}})^{p_2}\cdot A_{2n-3}^{\frac{t_1}{2}}\}^{p_3}A_{2n-4}^{-\frac{t_2}{2}}]^{p_4}A_{2n-5}^{\frac{t_2}{2}}\cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$;

(II.2n-1) $A_1^{r-t_n} \leqslant \Big\{A_1^{\frac{r}{2}}\Big[A_1^{-\frac{t_n}{2}}\{A_2^{\frac{t_{n-1}}{2}}\cdots A_{2n-4}^{\frac{t_2}{2}}[A_{2n-3}^{-\frac{t_2}{2}}\{A_{2n-2}^{\frac{t_1}{2}}\cdot(A_{2n-1}^{-\frac{t_1}{2}}A_{2n}^{p_1}A_{2n-1}^{-\frac{t_1}{2}})^{p_2}\cdot A_{2n-2}^{\frac{t_1}{2}}\}^{p_3}A_{2n-3}^{-\frac{t_2}{2}}]^{p_4}A_{2n-4}^{\frac{t_2}{2}}\cdots A_2^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}}A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}A_1^{\frac{r}{2}}\Big\}^{\frac{1}{m}}$.

(III) There exists strictly positive operators $S_1, S_2, S_3, \cdots, S_{2n-2}, S_{2n-1}$ satisfying the following operator equalities, respectively, where each S_i $(i = 1, 2, \cdots, 2n-1)$ is unique with $\|S_i\| \leqslant 1$.

(III.1) $A_{2n}^{-\frac{t_n}{2}} S_1 (A_{2n}^{r-t_n} S_1)^{m-1} A_{2n}^{-\frac{t_n}{2}} = A_{2n}^{-\frac{t_n}{2}} (S_1 A_{2n}^{r-t_n})^{m-1} S_1 A_{2n}^{-\frac{t_n}{2}} =$
$\Big[A_{2n}^{-\frac{t_n}{2}} \{A_{2n-1}^{\frac{t_{n-1}}{2}} \cdots A_5^{\frac{t_2}{2}} \big[A_4^{-\frac{t_2}{2}} \{A_3^{\frac{t_1}{2}} (A_2^{-\frac{t_1}{2}} A_1^{p_1} A_2^{-\frac{t_1}{2}})^{p_2} A_3^{\frac{t_1}{2}}\}^{p_3} A_4^{-\frac{t_2}{2}}\big]^{p_4} A_5^{\frac{t_2}{2}} \cdots A_{2n-1}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}} A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}$;

(III.2) $A_{2n}^{-\frac{t_n}{2}} S_2 (A_{2n}^{r-t_n} S_2)^{m-1} A_{2n}^{-\frac{t_n}{2}} = A_{2n}^{-\frac{t_n}{2}} (S_2 A_{2n}^{r-t_n})^{m-1} S_2 A_{2n}^{-\frac{t_n}{2}} =$
$\Big[A_{2n}^{-\frac{t_n}{2}} \{A_{2n}^{\frac{t_{n-1}}{2}} \cdots A_6^{\frac{t_2}{2}} \big[A_5^{-\frac{t_2}{2}} \{A_4^{\frac{t_1}{2}} (A_3^{-\frac{t_1}{2}} A_2^{p_1} A_3^{-\frac{t_1}{2}})^{p_2} A_4^{\frac{t_1}{2}}\}^{p_3} A_5^{-\frac{t_2}{2}}\big]^{p_4} A_6^{\frac{t_2}{2}} \cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}} A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}$;

(III.3) $A_{2n}^{-\frac{t_n}{2}} S_3 (A_{2n}^{r-t_n} S_3)^{m-1} A_{2n}^{-\frac{t_n}{2}} = A_{2n}^{-\frac{t_n}{2}} (S_3 A_{2n}^{r-t_n})^{m-1} S_3 A_{2n}^{-\frac{t_n}{2}} =$
$\Big[A_{2n}^{-\frac{t_n}{2}} \{A_{2n}^{\frac{t_{n-1}}{2}} \cdots A_7^{\frac{t_2}{2}} \big[A_6^{-\frac{t_2}{2}} \{A_5^{\frac{t_1}{2}} (A_4^{-\frac{t_1}{2}} A_3^{p_1} A_4^{-\frac{t_1}{2}})^{p_2} A_5^{\frac{t_1}{2}}\}^{p_3} A_6^{-\frac{t_2}{2}}\big]^{p_4} A_7^{\frac{t_2}{2}} \cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}} A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}$;

$\cdots\cdots\cdots\cdots$

(III.n) $A_{2n}^{-\frac{t_n}{2}} S_n (A_{2n}^{r-t_n} S_n)^{m-1} A_{2n}^{-\frac{t_n}{2}} = A_{2n}^{-\frac{t_n}{2}} (S_n A_{2n}^{r-t_n})^{m-1} S_n A_{2n}^{-\frac{t_n}{2}} =$
$\Big[A_{2n}^{-\frac{t_n}{2}} \{A_{2n}^{\frac{t_{n-1}}{2}} \cdots A_{n+4}^{\frac{t_2}{2}} \big[A_{n+3}^{-\frac{t_2}{2}} \{A_{n+2}^{\frac{t_1}{2}} (A_{n+1}^{-\frac{t_1}{2}} A_n^{p_1} A_{n+1}^{-\frac{t_1}{2}})^{p_2} A_{n+2}^{\frac{t_1}{2}}\}^{p_3} A_{n+3}^{-\frac{t_2}{2}}\big]^{p_4} A_{n+4}^{\frac{t_2}{2}} \cdots A_{2n}^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}} A_{2n}^{-\frac{t_n}{2}}\Big]^{p_{2n}}$;

(III.n+1) $A_1^{-\frac{t_n}{2}} S_{n+1}^{-1} (A_1^{r-t_n} S_{n+1}^{-1})^{m-1} A_1^{-\frac{t_n}{2}} = A_1^{-\frac{t_n}{2}} (S_{n+1}^{-1} A_1^{r-t_n})^{m-1} S_{n+1}^{-1} A_1^{-\frac{t_n}{2}} =$
$\Big[A_1^{-\frac{t_n}{2}} \{A_1^{\frac{t_{n-1}}{2}} \cdots A_{n-2}^{\frac{t_2}{2}} \big[A_{n-1}^{-\frac{t_2}{2}} \{A_n^{\frac{t_1}{2}} (A_{n+1}^{-\frac{t_1}{2}} A_{n+2}^{p_1} A_{n+1}^{-\frac{t_1}{2}})^{p_2} A_n^{\frac{t_1}{2}}\}^{p_3} A_{n-1}^{-\frac{t_2}{2}}\big]^{p_4} A_{n-2}^{\frac{t_2}{2}} \cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}} A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}$;

$\cdots\cdots\cdots\cdots$

(III.2n-2) $A_1^{-\frac{t_n}{2}} S_{2n-2}^{-1} (A_1^{r-t_n} S_{2n-2}^{-1})^{m-1} A_1^{-\frac{t_n}{2}} = A_1^{-\frac{t_n}{2}} (S_{2n-2}^{-1} A_1^{r-t_n})^{m-1} S_{2n-2}^{-1} A_1^{-\frac{t_n}{2}} =$
$\Big[A_1^{-\frac{t_n}{2}} \{A_1^{\frac{t_{n-1}}{2}} \cdots \big[A_{2n-4}^{-\frac{t_2}{2}} \{A_{2n-3}^{\frac{t_1}{2}} (A_{2n-2}^{-\frac{t_1}{2}} A_{2n-1}^{p_1} A_{2n-2}^{-\frac{t_1}{2}})^{p_2} A_{2n-3}^{\frac{t_1}{2}}\}^{p_3} A_{2n-4}^{-\frac{t_2}{2}}\big]^{p_4} \cdots A_1^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}} A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}$;

(III.2n-1) $A_1^{-\frac{t_n}{2}} S_{2n-1}^{-1} (A_1^{r-t_n} S_{2n-1}^{-1})^{m-1} A_1^{-\frac{t_n}{2}} = A_1^{-\frac{t_n}{2}} (S_{2n-1}^{-1} A_1^{r-t_n})^{m-1} S_{2n-1}^{-1} A_1^{-\frac{t_n}{2}} =$
$\Big[A_1^{-\frac{t_n}{2}} \{A_2^{\frac{t_{n-1}}{2}} \cdots \big[A_{2n-3}^{-\frac{t_2}{2}} \{A_{2n-2}^{\frac{t_1}{2}} (A_{2n-1}^{-\frac{t_1}{2}} A_{2n}^{p_1} A_{2n-1}^{-\frac{t_1}{2}})^{p_2} A_{2n-2}^{\frac{t_1}{2}}\}^{p_3} A_{2n-3}^{-\frac{t_2}{2}}\big]^{p_4} \cdots A_2^{\frac{t_{n-1}}{2}}\}^{p_{2n-1}} A_1^{-\frac{t_n}{2}}\Big]^{p_{2n}}$;

Proof. Let $A_{2n+1} = A_{2n}$ in Theorem 3.1. □

Together with Theorem 3.1 and Theorem 3.2, we give an application of the characterizations of $A_k \geqslant A_{k-1} \geqslant \cdots \geqslant A_2 \geqslant A_1 > 0$ to operator equalities for any positive integer k.

References

[1] A. Aluthge, On p-hyponormal operators for $0 < 0 < 1$, Integr. Equat. Oper. Th., **13** (1990), 307-315.

[2] A. Aluthge and D. Wang, Powers of p-hyponormal operators, J. Inequal. Appl., **3** (1999), 279-284.

[3] R. G. Douglas, On majorization, factorization, and range inclusion of operators on Hilbert space, Proc. Amer. Math. Soc., **17** (1966), 413-415.

[4] M. Fujii, T. Furuta and E. Kamei, Furuta's inequality and its applications to Ando's theorem, Linear Algebra Appl., **179** (1993), 161-169.

[5] M.Fujii, S. Izumino, R. Nakamoto and Y. Seo, Operator inequalities related to Cauchy-Schwarz and Hölder-McCarthy inequalities, Nihonkai Math. J., **8** (1997), 117-122.

[6] M. Fujii, A. Matsumoto and R. Nakamoto, A short proof of the best possibility for the grand Furuta inequality, J. Inequal. Appl., **4** (1999), 339-344.

[7] M. Fujii, E. Kamei and R. Nakamoto, On a question of Furuta on chaotic order, Linear Algebra Appl., **341** (2002), 119-127.

[8] T. Furuta, $A \geqslant B \geqslant 0$ assures $(B^rA^pA^r)^{1/q} \geqslant B^{\frac{p+2r}{q}}$ for $r \geqslant 0$, $p \geqslant 0$, $q \geqslant 1$ with $(1+2r)q \geqslant p+2r$, Proc. Amer. Math. Soc., **101** (1987), 85-88.

[9] T. Furuta, Furuta's inequality and its application to the realtive operator entropy, J. Operator Theory, **30** (1993), 21-30.

[10] T. Furuta, An extension of the Furuta inequality and Ando-Hiai log majorization, Linear Algebra Appl., **219** (1995), 139-155.

[11] T. Furuta, Further extension of an order preserving operator inequality, J. Math. Inequal., **2**, 4 (2008), 465-472.

[12] T. Huruya, A note on p-hyponormal operators, Proc. Amer. Math. Soc., **125** (1997), 3617-3624.

[13] M. Ito and T. Yamazaki, Relations between two inequalities $(B^{\frac{r}{2}}A^pB^{\frac{r}{2}})^{\frac{r}{p+r}} \geqslant B^r$ and $(A^{\frac{p}{2}}B^rA^{\frac{p}{2}}) \leqslant A^p$ and their applications, Integr. Equat. Oper. Th., **44** (2002), 442-450.

[14] C.-S. Lin, On operator order and chaotic operator order for two operators, Linear Algebra Appl., **425** (2007), 1-6.

[15] C.-S. Lin and Y. J. Cho, Characterizations of the operator inequality $A \geqslant B \geqslant C$, Math. Inequal. Appl., **14**, 3 (2011), 575-580.

[16] K. Tanahashi, Best possibility of Furuta inequality, Proc. Amer. Math. Soc., **124** (1996), 141-146.

[17] K. Tanahashi, The best possibility of the grand Furuta inequality, Proc. Amer. Math. Soc., **128** (2000), 511-519.

[18] M. Uchiyama, Criteria for monotonicity of operator means, J. Math. Soc. Japan, **55** (2003), 197-207.

[19] T. Yamazaki, Simplified proof of Tanahashi's result on the best possibility of generalized Furuta inequality, Math. Inequal. Appl., **2**, 3 (1999), 473-477.

[20] C. Yang and Y. Wang, Further extension of Furuta inequality, J. Math. Inequal., **4**, 3 (2010), 391-398.

[21] J. Yuan and Z. Gao, Classified construction of generalized Furuta type operator functions, Math. Inequal. Appl., **11**, 2 (2008), 189-202.

[22] J. Yuan, Classified construction of generalized Furuta type operator functions, Math. Inequal. Appl., **13**, 4 (2010), 775-784.

[23] J. Yuan, Furuta inequality and q-hyponormal operators, Oper. Matrices, **4**, 3 (2010), 405-415.